

AD-A228 293

DTIC FILE COPY

ARJ 25 839.33-MA

2

ESTIMATING THE ASYMPTOTIC VARIANCE
WITH BATCH MEANS

by

Peter W. Glynn
Ward Whitt

TECHNICAL REPORT No. 57

March 1990

Prepared under the Auspices
of

U.S. Army Research Contract
DAAL-03-88-K-0063

DTIC
ELECTE
OCT 26 1990
S B D
CD

Approved for public release: distribution unlimited.

Reproduction in whole or in part is permitted for any
purpose of the United States government.

DEPARTMENT OF OPERATIONS RESEARCH
STANFORD UNIVERSITY
STANFORD, CALIFORNIA

90 10 28 095

ESTIMATING THE ASYMPTOTIC VARIANCE
WITH BATCH MEANS

by

Peter W. Glynn*

Department of Operations Research
Stanford University
Stanford, CA 94305-4022

and

Ward Whitt

AT&T Bell Laboratories
Murray Hill, NJ 07974-2070

Abstract

We show that the asymptotic variance constant in a stochastic simulation cannot be estimated consistently from batch means when the number of batches is held fixed as the run length increases.

Keywords: simulation, estimation, batch means, consistency.

*Research supported by the U.S. Army Research Office under Contract DAAL-03-88-K-0063.

1. Introduction

In this paper ~~we show~~ that there does not exist a procedure to consistently estimate the asymptotic variance constant in a stochastic simulation using batch means when the number of batches is held fixed as the run length increases. Thus, if consistency is desired, then the number of batches must increase as the run length increases.

To be precise, we must first specify what we mean by an estimation procedure. To be interesting, an estimation procedure should apply to a large family of stochastic processes. Hence, let $X \equiv \{X(t) : t \geq 0\}$ be a measurable mapping from a measure space (Ω, \mathcal{F}) into $D \equiv D[0, \infty)$, the space of right-continuous real-valued functions on the interval $[0, \infty)$ with left limits, endowed with the usual Skorohod topology and associated Borel σ -field; e.g., see Ethier and Kurtz [3]. Of course, we want the underlying space (Ω, \mathcal{F}) to be sufficiently rich; it suffices to let $\Omega = D$ and $X(t)$ be the projection or coordinate map. We consider the set \mathcal{P} of all probability measures P on (Ω, \mathcal{F}) such that there exist finite deterministic constants $\mu \equiv \mu(P)$ and $\sigma \equiv \sigma(P)$ such that

$$n^{1/2} \left(\frac{1}{n} \int_0^{[n]} \dot{X}(s) - \mu \right) \Rightarrow \sigma B(t) \text{ as } n \rightarrow \infty, \quad (1)$$

where \Rightarrow denotes weak convergence in D with respect to P and $B \equiv \{B(t) : t \geq 0\}$ is standard (zero-drift, unit diffusion coefficient) Brownian motion. Our goal is to estimate σ^2 , but we want our procedure to apply to all $P \in \mathcal{P}$. In other words, the procedure should apply to all stochastic processes X in D satisfying the functional central limit theorem (FCLT) (1).

To apply the method of batch means, we specify the number m of batches and the total run length T . We then construct our estimates from the m non-overlapping intervals of length T/m ; i.e., let the i^{th} batch mean be

$$\bar{X}_i(T) = \frac{m}{T} \int_{(i-1)T/M}^{iT/M} X(s) ds, \quad 1 \leq i \leq m. \quad (2)$$

We now want a procedure for combining the m observations $\bar{X}_1(T), \dots, \bar{X}_m(T)$ in such a way that σ^2 is consistently estimated as $T \rightarrow \infty$. This "combining transformation" should not depend on the "fine structure" of the process X . In particular, it should not depend on μ and σ^2 . Thus, in this context we say that an *estimation procedure* is a family of measurable mappings

$$g_T : R^m \rightarrow R \quad \text{for } T > 0, \quad (3)$$

such that the estimate of σ^2 is $g_T(x_1, \dots, x_m)$ when the total run length is T and $\bar{X}_i(t) = x_i$, $1 \leq i \leq m$. Note that g_T can depend on T , but is independent of P .

We say that an estimation procedure is \mathcal{P} -consistent if for each $P \in \mathcal{P}$

$$g_T(\bar{X}_1(T), \dots, \bar{X}_m(T)) \Rightarrow \sigma^2(P) \text{ as } T \rightarrow \infty. \quad (4)$$

Here \Rightarrow denotes weak convergence with respect to P in R , which is equivalent to convergence in probability since $\sigma^2(P)$ is deterministic. Since we have a negative result, we focus on this weak consistency. We would have strong consistency if the convergence was w.p.1 with respect to P .

Here is our main result. It applies to any m .

Theorem 1. There does not exist an estimation procedure that is \mathcal{P} -consistent.

In Section 2 we show what happens with the standard variance estimator. We see that we do not get consistency for σ^2 for any fixed m , but we can get as close as we wish by letting m be suitably large. In Section 3 we prove Theorem 1.

Theorem 1 has applications to sequential stopping. It shows that the sufficient conditions in Glynn and Whitt [5] for asymptotic validity are *not satisfied* when the number of batches is held fixed as the run length grows.

For a fixed run length, our analysis shows that it is desirable to pick the number of batches as large as possible without seriously violating the assumption that the batches are independent and identically distributed (i.i.d.) with a normal distribution. However, the i.i.d. normal assumption typically holds only as an approximation and then only when there are large batch sizes. Statistical tests can be used to validate the assumption, but repeated tests of significance on the same data are fraught with peril, both theoretically and empirically. Hence, Schmeiser [9] suggested using a relatively small fixed number of batches, e.g., about 20. This avoids the complications above and gives relatively robust confidence intervals. However, we show that this is achieved at the expense of consistency for the variance estimator. Asymptotically valid confidence intervals are obtained anyway of course by cancellation methods, i.e., using the t distribution. For further discussion, see Schmeiser [9], Goldsman and Meketon [6], Sargent, Kang and Goldsman [8], Glynn and Iglehart [4] and Damerdjani [2].

2. The Standard Estimator

The standard estimation procedure is specified by

$$g_T^*(x_1, \dots, x_m) = \frac{T}{m(m-1)} \sum_{i=1}^m \left[x_i - \frac{1}{m} \sum_{k=1}^m x_k \right]^2 \quad (5)$$

for all $T > 0$, $m \geq 2$ and $(x_1, \dots, x_m) \in R^m$. Let $\stackrel{d}{=}$ denote equality in distribution.

Theorem 2. Under (1),

$$g_T^*(\bar{X}_1(T), \dots, \bar{X}_m(T)) \Rightarrow \sigma^2 m^2 g_1^*([B(i/m) - B((i-1)/m)], 1 \leq i \leq m) \\ = \frac{\sigma^2 \chi_{m-1}^2}{m-1} \text{ in } R,$$

where χ_{m-1}^2 is a chi-square random variable with $m-1$ degrees of freedom.

Proof. Note that

Distribution For	
GRA&I	<input checked="" type="checkbox"/>
TAB	<input type="checkbox"/>
unched	<input type="checkbox"/>
ocation	

Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



$$g_T^*(\bar{X}_1(T), \dots, \bar{X}_m(T)) = \frac{m}{m-1} \sum_{i=1}^m \left[\sqrt{T} \frac{\int_{(i-1)T/m}^{iT/m} X(s) ds}{T} - \sqrt{T} \frac{\int_0^T X(s) ds}{mT} \right]^2$$

$$\Rightarrow \sigma^2 \frac{m}{m-1} \sum_{i=1}^m \left[B(i/m) - B((i-1)/m) - \frac{B(1)}{m} \right]^2 \stackrel{d}{=} \frac{\sigma^2 \chi_{m-1}^2}{m-1} \text{ by (1)} \quad \blacksquare$$

Note that $\sigma^2 \chi_{m-1}^2 / m-1$ has mean σ^2 and variance $2\sigma^4 / (m-1)$; see p. 168 of Johnson and Kotz [7]. Moreover, as m increases,

$$\frac{\sigma^2 \chi_m^2}{m} \Rightarrow \sigma^2 \quad (6)$$

and

$$\sqrt{m} \left[\frac{\sigma^2 \chi_m^2}{m} - \sigma^2 \right] \Rightarrow N(0, 2\sigma^4) \quad (7)$$

where $N(m, \sigma^2)$ denotes a normally distributed random variable with mean m and variance σ^2 . Hence, we can get as close as we want if we choose m suitably large. Moreover, we can obtain consistency under extra regularity conditions if $m \rightarrow \infty$ and $T \rightarrow \infty$ so that $T/m \rightarrow \infty$; see Goldsman and Meketon [6] and Damerdjy [2]. In fact, Damerdjy even proves strong consistency for a class of stochastic processes.

3. Proof of Theorem 1

To establish the negative result, it suffices to restrict attention to probability measures P such that X coincides with σB where B is standard Brownian motion. Then, for any m and T ,

$$(\bar{X}_1(T), \dots, \bar{X}_m(T)) = \left[\frac{m}{T} \sigma B(T/m), \dots, \frac{m}{T} \sigma [B(T) - B(T(m-1)/m)] \right], \quad (7)$$

so that the batch means are distributed exactly as m i.i.d. normal random variables with mean 0 and variance $\sigma^2 m/T$. Without loss of generality, we can remove the m/T factor by

considering the transformed functions

$$\tilde{g}_T(x_1, \dots, x_m) = g_T \left[\sqrt{\frac{T}{m}} x_1, \dots, \sqrt{\frac{T}{m}} x_m \right]. \quad (8)$$

Note that

$$g_T(\bar{X}_1(T), \dots, \bar{X}_m(T)) \stackrel{d}{=} \tilde{g}_T(\sigma N) \quad \text{for all } T, \quad (9)$$

where $\sigma N \equiv (\sigma N_1, \dots, \sigma N_m)$ and N is a fixed vector of i.i.d. standard (mean 0, variance 1) normal random variables.

Now consistency requires that

$$\tilde{g}_T(\sigma N) \Rightarrow \sigma \text{ as } T \rightarrow \infty \quad (10)$$

for all $\sigma \geq 0$, but this cannot happen for two or more different positive values of σ , say σ_1 and σ_2 . To see this, first note that the convergence in probability for σ_1 in (10) implies that there is a sequence $\{T_n : n \geq 1\}$ of deterministic positive numbers with $T_n \rightarrow \infty$ such that

$$\tilde{g}_{T_n}(\sigma_1 N) \rightarrow \sigma_1 \quad \text{w.p.1 as } n \rightarrow \infty; \quad (11)$$

see Theorem 4.2.3 of Chung [1]. By (10), $\tilde{g}_{T_n}(\sigma_2 N) \Rightarrow \sigma_2$ as $n \rightarrow \infty$. Hence, there is a deterministic subsequence $\{T'_n : n \geq 1\}$ of $\{T_n : n \geq 1\}$ such that

$$\tilde{g}_{T'_n}(\sigma_i N) \rightarrow \sigma_i \quad \text{w.p.1 as } n \rightarrow \infty \quad (12)$$

for both $i = 1$ and 2 . Hence, for $i = 1$ and 2 , $\tilde{g}_{T'_n}(x) \rightarrow \sigma_i$ for almost all x with respect to the law of $\sigma_i N$, which implies that $\tilde{g}_{T'_n}(x) \rightarrow \sigma_i$ for almost all x with respect to Lebesgue measure on R^m , since $\sigma_i N$ has a positive density with respect to Lebesgue measure. However, it is not possible to have $\tilde{g}_{T'_n}(x)$ simultaneously converge almost everywhere with respect to Lebesgue measure to two different limits. (The set of convergence to one limit must be contained in the null set of non-convergence for the other limit.)

References

- [1] Chung, K. L. (1974) *A Course in Probability Theory*, Second ed., Academic Press, New York.
- [2] Damerджи, H. (1989) Strong consistency of the variance estimator in steady-state simulation output analysis. Department of Operations Research, Stanford University.
- [3] Ethier, S. N. and T. G. Kurtz (1986) *Markov Processes, Characterization and Convergence*, Wiley, New York.
- [4] Glynn, P. W. and D. L. Iglehart (1990) Simulation output analysis using standardized times series. *Math. Oper. Res.* 15, 1-16.
- [5] Glynn, P. W. and W. Whitt (1989) The asymptotic validity of sequential stopping rules in stochastic simulation. Submitted for publication.
- [6] Goldsman, D. and M. Meketon (1986) A comparison of several variance estimators. School of Industrial and Systems Engineering, Georgia Institute of Technology.
- [7] Johnson, N. L. and S. Kotz (1970) *Continuous Univariate Distributions-1, Distributions in Statistics*, Wiley, New York.
- [8] Sargent, R. G., K. Kang and D. Goldsman (1989) An investigation of finite sample behavior of confidence interval estimation procedures. Department of Industrial Engineering, Syracuse University.
- [9] Schmeiser, B. (1982) Batch size effects in the analysis of simulation output. *Oper. Res.* 30, 556-568.

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S) AR 25839.33-MA	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 57		7a. NAME OF MONITORING ORGANIZATION U. S. Army Research Office	
6a. NAME OF PERFORMING ORGANIZATION Dept. of Operations Research	6b. OFFICE SYMBOL (If applicable)	7b. ADDRESS (City, State, and ZIP Code) P. O. Box 12211 Research Triangle Park, NC 27709-2211	
6c. ADDRESS (City, State, and ZIP Code) Stanford, CA 94305-4022	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAA03-88-K-0063		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U. S. Army Research Office	8b. OFFICE SYMBOL (If applicable)	10. SOURCE OF FUNDING NUMBERS	
8c. ADDRESS (City, State, and ZIP Code) P. O. Box 12211 Research Triangle Park, NC 27709-2211		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Estimating the asymptotic variance with batch means			
12. PERSONAL AUTHOR(S) Peter W. Glynn and Ward Whitt			
13a. TYPE OF REPORT Technical	13b. TIME COVERED FROM TO	14. DATE OF REPORT (Year, Month, Day) March 1990	15. PAGE COUNT 7
16. SUPPLEMENTARY NOTATION The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	simulation, estimation, batch means, consistency.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) We show that the asymptotic variance constant in a stochastic simulation cannot be estimated consistently from batch means when the number of batches is held fixed as the run length increases.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL		22b. TELEPHONE (Include Area Code)	22c. OFFICE SYMBOL